

DTP/98/5
March 1998

Parafermionic representation of the affine $\hat{sl}(2|1; \mathbb{C})$ algebra at fractional level.

P. Bowcock, M. Hayes and A. Taormina

Department of Mathematical Sciences, University of Durham, Durham, DH1 3LE, England

Abstract

The four fermionic currents of the affine superalgebra $\hat{sl}(2|1; \mathbb{C})$ at fractional level $k = \frac{1}{u} - 1, u \in \mathbb{N}$ are shown to be realised in terms of a free scalar field, an $\hat{sl}(2; \mathbb{C})$ doublet field and a primary field of the parafermionic algebra $\mathbb{Z}_{u-1}$.

e-mail: Peter.Bowcock@durham.ac.uk, M.R.Hayes@durham.ac.uk, Anne.Taormina@durham.ac.uk

Affine Lie algebras and superalgebras at fractional level have received some attention in the last years in the context of the study of non critical strings [1, 2, 3]. Our original motivation in the analysis of the representation theory of $\hat{sl}(2|1; \mathbb{C})$ at fractional level k is its potential relevance in the description of $N = 2$ (non critical) superstrings. We take the matter coupled to supergravity in an $N = 2$ super Coulomb gas representation with central charge

$$c_{\text{matter}} = 3(1 - \frac{2p}{u}), \quad p, u \in \mathbb{N}, \quad \gcd(p, u) = 1. \quad (1)$$

Then, the level k of the matter affine superalgebra $\hat{sl}(2|1)$ appearing in the $SL(2|1; \mathbb{R})/SL(2|1; \mathbb{R})$ gauged Wess-Zumino-Novikov-Witten model, believed to be intimately related to the $N = 2$ string, is of the form

$$k = \frac{p}{u} - 1. \quad (2)$$

This is precisely the type of fractional levels first discussed in the paper by Kac and Wakimoto on admissible representations of affine Lie algebras [4]. The detailed study of the representation theory of $\hat{sl}(2|1; \mathbb{C})$ for levels of the form (2) with $p = 1$ (which leads to the description of *unitary* minimal $N = 2$ matter) has revealed the existence of a finite number of irreducible representations whose characters form a finite representation of the modular group [5]. There are thus *rational*, although non unitary, theories associated with $\hat{sl}(2|1; \mathbb{C})$ at fractional level.

One of the very interesting byproducts of the above analysis is the existence of a representation of the $\hat{sl}(2|1; \mathbb{C})$ currents in terms of $\mathbb{Z}_{u-1}$ parafermions, as we shall now describe.

At level $k = \frac{1}{u} - 1$, the vacuum representation is in the Ramond sector of the theory and is labeled by the two quantum numbers $h_-^R = 0$ and $h_+^R = 0$, which correspond to the isospin and $U(1)$ charges of the highest weight state [6, 8]. The associated character is given by,

$$\chi_{0,0}^{R, IV, \hat{sl}(2|1; \mathbb{C})_k}(\tau, \sigma, \nu) = F^R(\tau, \sigma, \nu) \times \sum_{a \in \mathbb{Z}} q^{a^2 pu} z^{-ap} \frac{1 - q^{2ua} z^{-1}}{(1 + q^{au} z^{-\frac{1}{2}} \zeta^{-\frac{1}{2}})(1 + q^{au} z^{-\frac{1}{2}} \zeta^{\frac{1}{2}})}, \quad (3)$$

where $q = e^{2i\pi\tau}$, $z = e^{2i\pi\sigma}$, $\zeta = e^{2i\pi\nu}$ and the factor $F^R(\tau, \sigma, \nu)$ is,

$$F^R(\tau, \sigma, \nu) = \prod_{n=1}^{\infty} \frac{(1 + z^{\frac{1}{2}} \zeta^{\frac{1}{2}} q^n)(1 + z^{-\frac{1}{2}} \zeta^{\frac{1}{2}} q^{n-1})(1 + z^{\frac{1}{2}} \zeta^{-\frac{1}{2}} q^n)(1 + z^{-\frac{1}{2}} \zeta^{-\frac{1}{2}} q^{n-1})}{(1 - q^n)^2 (1 - z q^n)(1 - z^{-1} q^{n-1})}. \quad (4)$$

The crucial ingredient needed for the construction of the new representation of $\hat{sl}(2|1; \mathbb{C})$ currents is provided by the decomposition of the above character in characters of the affine $\hat{sl}(2; \mathbb{C})$ subalgebra at the same fractional level. The general decomposition formulae appear in [5] for the Neveu-Schwarz sector of the theory. Using spectral flow as discussed in [8], and the following relation between the $\hat{sl}(2; \mathbb{C})$ characters of isospin $j(n, n') = \frac{1}{2}(n - n'(k + 2))$ and isospin $j(u - n - 1, u - n' - 1)$,

$$\chi_{n, n'}^{\hat{sl}(2; \mathbb{C})_k}(\tau, -\sigma - \tau) = z^{-\frac{k}{2}} q^{-\frac{k}{4}} \chi_{u-n-1, u-n'-1}^{\hat{sl}(2; \mathbb{C})_k}(\tau, \sigma), \quad (5)$$

one easily obtains the Ramond vacuum decomposition,

$$\begin{aligned} \chi_{0,0}^{R,IV,\hat{sl}(2|1;\mathbb{C})_k}(\tau, \sigma, \nu) = \\ \sum_{i=0}^{u-1} \sum_{s=0}^{u-2} c_{i,-3i-2us}^{(u-1)}(\tau) \theta_{u(u-1)(i+1)+2iu(\frac{u}{2}-[\frac{u}{2}])-2us,u(u-1)}(\tau, \frac{\nu}{u}) \chi_{u-1-i,0}^{\hat{sl}(2;\mathbb{C})_k}(\tau, \sigma), \end{aligned} \quad (6)$$

where the symbol $[\frac{u}{2}]$ is the integer part of $\frac{u}{2}$. The above formula provides some insight into the coset structure $\frac{\hat{sl}(2|1;\mathbb{C})_k}{sl(2;\mathbb{C})_k}$. Indeed, the functions $c_{\ell,m}^{(u-1)}(\tau)$ are the $\hat{su}(2)_{u-1}$ string functions introduced in [9]. When multiplied by the Dedekind function $\eta(\tau)$, they provide the partition functions of the parafermionic algebra $\mathbb{Z}_{u-1}$ whose lowest dimensional fields are labeled Φ_m^ℓ [10, 11, 12]. Furthermore, the theta functions at level $u(u-1)$, when divided by the Dedekind function, are the characters for the rational torus $A_{u(u-1)}$ [13]. The coset structure is also reflected in the following relations between central charges. The central charge of the Virasoro algebra associated to $\hat{sl}(2|1;\mathbb{C})_k$ is,

$$c_{\hat{sl}(2/1)} = \frac{k \text{ sdim } sl(2/1)}{k+1} = 0 \quad (7)$$

since there is an equal number of bosonic and fermionic generators in $sl(2/1)$. As a consequence, the central charge of the coset $\frac{\hat{sl}(2|1;\mathbb{C})_k}{sl(2;\mathbb{C})_k}$, for $k = \frac{1}{u} - 1$, is

$$c_{\text{coset}} = -c_{sl(2)} = \frac{3(u-1)}{u+1}. \quad (8)$$

It can in turn be rewritten as,

$$c_{\text{coset}} = c_{\text{torus}} + c_{\text{paraf}} = 1 + \frac{2(u-2)}{u+1}. \quad (9)$$

A close reading of the decomposition formula (6) provides the information needed to construct the new $\hat{sl}(2|1;\mathbb{C})_k$ representation. By expanding the vacuum character (3) in powers of q , one discovers at grade one the eight states belonging to the adjoint representation of $sl(2/1;\mathbb{C})$. It is also straightforward to identify on the right hand side of the decomposition (6), the four states at grade one which are neutral with respect to the $U(1)$ algebra, i.e. those which are independent of the variable ζ . They all lie in the term $i = u-1$, $s = 0$ and correspond to the first bosonic excitations of the vacuum. The other four states correspond to the first fermionic vacuum excitations and lie in the terms $i = u-2$, $s = [\frac{u}{2}]$, $s = [\frac{u}{2}] - 1$. The string function appearing in these two terms is $c_{1,1}^{(u-1)}(\tau)$, which is, upon multiplication by the Dedekind function, the parafermionic partition function with lowest dimensional field $\Phi_1^1(z)$, identified with the primary field $\sigma_{(1)}(z)$ with conformal dimension [10],

$$h_{(1)} = \frac{u-2}{2(u-1)(u+1)}. \quad (10)$$

This field is one of the $(u-2)$ conformal spin fields or order parameters associated with the parafermion algebra based on $\mathbb{Z}_{u-1}$ [10]. It enters in our representation of the fermionic

$\hat{sl}(2|1; \mathbb{C})$ currents as follows. Adopting the notations introduced in [6] for the $\hat{sl}(2|1; \mathbb{C})$ currents, one writes,

$$\begin{aligned} j^\pm(z) &= \Delta^\pm(z) \left(i \sqrt{\frac{1-u}{u}} \right) : e^{\mp \frac{i}{2} \sqrt{\frac{2u}{u-1}} \phi(z)} : \sigma_{(1)}(z), \\ j^{\pm'}(z) &= \Delta^\pm(z) \left(i \sqrt{\frac{1-u}{u}} \right) : e^{\pm \frac{i}{2} \sqrt{\frac{2u}{u-1}} \phi(z)} : \sigma_{(1)}(z), \end{aligned} \quad (11)$$

where $\Delta^\pm(z)$ is the field corresponding to the $\hat{sl}(2; \mathbb{C})_k$ doublet highest weight state with conformal weight $h_{\text{doublet}} = \frac{3u}{4(u+1)}$ and O.P.E. [14],

$$\begin{aligned} \Delta^+(z) \Delta^-(w) &= (z-w)^{-\frac{3u}{2(u+1)}} + \frac{u}{1-u} (z-w)^{-\frac{u-2}{2(u+1)}} J_3(w) + O((z-w)^{\frac{u}{2(u+1)}}), \\ \Delta^\pm(z) \Delta^\pm(w) &= \frac{u}{1-u} (z-w)^{-\frac{u-2}{2(u+1)}} J^\pm(w) + O((z-w)^{\frac{u}{2(u+1)}}), \end{aligned} \quad (12)$$

where $J^\pm(w)$ and $J^3(w)$ are the $\hat{sl}(2; \mathbb{C})$ currents. As usual, the vertex operators have conformal weight $h_{U(1)} = \frac{u}{4(u-1)}$ and obey the O.P.E.,

$$: e^{\pm \frac{i}{2} \sqrt{\frac{2u}{u-1}} \phi(z)} :: e^{\pm \frac{i}{2} \sqrt{\frac{2u}{u-1}} \phi(w)} : = (z-w)^{\frac{u}{2(u-1)}} : e^{\pm \frac{i}{2} \sqrt{\frac{2u}{u-1}} (\phi(z) + \phi(w))} : \quad (13)$$

and

$$: e^{\pm \frac{i}{2} \sqrt{\frac{2u}{u-1}} \phi(z)} :: e^{\mp \frac{i}{2} \sqrt{\frac{2u}{1-u}} \phi(w)} : = (z-w)^{-\frac{u}{2(u-1)}} : e^{\pm \frac{i}{2} \sqrt{\frac{2u}{u-1}} (\phi(z) - \phi(w))} : . \quad (14)$$

Note that

$$h_{\text{doublet}} + h_{U(1)} + h_{(1)} = 1, \quad (15)$$

as required for the fermionic $\hat{sl}(2|1; \mathbb{C})_k$ currents. So, the parafermionic representation of $\hat{sl}(2|1; \mathbb{C})$ we obtained is in terms of the $\hat{sl}(2; \mathbb{C})$ currents, the $U(1)$ current

$$U(z) = i \sqrt{\frac{u-1}{2u}} \partial \phi(z), \quad (16)$$

where $\phi(z)$ is a free scalar field with the standard O.P.E.

$$\phi(z) \phi(w) \sim -\ln|z-w|, \quad (17)$$

and the four fermionic currents (11). Note that the parafermions enter in this representation in a rather different way than in representations of $\hat{sl}(2; \mathbb{C})$ at *integer* level k (see e.g. [15]), where the parafermionic algebra is $\mathbb{Z}_k$.

We now illustrate the case where $u = 3$, when the parafermions are the usual fermions. The spin field of interest is then $\sigma(z)$ of weight $1/16$, and appears in the description of the Ising model. The relevant O.P.E. are,

$$\begin{aligned} \sigma(z) \sigma(w) &= (z-w)^{-1/8} + \frac{1}{2} (z-w)^{3/8} \psi(w) + \dots \\ \Delta^+(z) \Delta^-(w) &= (z-w)^{-9/8} - \frac{3}{2} (z-w)^{-1/8} J^3(w) + \alpha (z-w)^{3/8} A_3(w) + \dots \\ \Delta^\pm(z) \Delta^\pm(w) &= -\frac{3}{2} (z-w)^{-1/8} J^\pm(w) + \alpha' (z-w)^{3/8} A^\pm(w) + \dots, \end{aligned} \quad (18)$$

where $\psi(w)$ is a fermion and $A^\pm(w), A_3(w)$ form an $\hat{sl}(2; \mathbb{C})$ triplet of weight $\frac{3}{2}$. These fields, whose occurrence in the theory can be traced back to the vacuum character formula (6), have to be taken into account in parafermionic representations of normal ordered products of currents, but are not relevant to the discussion of commutation relations of those currents. The exact value of the two parameters α and α' in the above O.P.E. is therefore not required for our purpose. Now, the short distance behaviour of $\hat{sl}(2|1; \mathbb{C})_k$ currents is given by,

$$\begin{aligned}
j^+(z)j^-(w) &\sim [U(w) + J^3(w)](z-w)^{-1} + k(z-w)^{-2}, \\
j^{+'}(z)j^{-'}(w) &\sim [U(w) - J^3(w)](z-w)^{-1} - k(z-w)^{-2}, \\
j^{\pm'}(z)j^{\pm}(w) &\sim J^\pm(w)(z-w)^{-1}, \\
U(z)U(w) &\sim -\frac{k}{2}(z-w)^{-2}, \quad J^3(z)J^3(w) \sim \frac{k}{2}(z-w)^{-2}, \\
2J^3(z)j^\pm(w) &\sim \pm j^\pm(w)(z-w)^{-1}, \quad 2J^3(z)j^{\pm'}(w) \sim \pm j^{\pm'}(w)(z-w)^{-1}, \\
2U(z)j^\pm(w) &\sim \mp j^\pm(w)(z-w)^{-1}, \quad 2U(z)j^{\pm'}(w) \sim \pm j^{\pm'}(w)(z-w)^{-1}, \\
J^\pm(z)j^\mp(w) &\sim \mp j^{\mp'}(w)(z-w)^{-1}, \quad J^\pm(z)j^{\mp'}(w) \sim \pm j^\pm(w)(z-w)^{-1}, \\
J^+(z)J^-(w) &\sim 2J^3(w)(z-w)^{-1} + k(z-w)^{-2}.
\end{aligned} \tag{19}$$

Using (13) and (14) together with the O.P.E. (18) truncated to the terms with negative powers of $(z-w)$, one checks that the fermionic currents as given in (11) and the bosonic current (16) provide, with the $\hat{sl}(2; \mathbb{C})$ currents $J^\pm(z)$ and $J^3(z)$, a representation of $\hat{sl}(2|1; \mathbb{C})$ at level $k = -2/3$.

A similar representation in the case $u = 4$ involves the spin field of the 3-states Potts model, with conformal weight $1/15$. This is hardly surprising, since the $\mathbb{Z}_2$ and $\mathbb{Z}_3$ parafermionic theories coincide with the Ising and 3-states Potts models respectively. At the level of characters, one actually has,

$$\chi_{1,1}^{Vir(3)}(\tau) = \eta(\tau)c_{2,2}^{(2)}(\tau), \quad \chi_{2,1}^{Vir(3)}(\tau) = \eta(\tau)c_{0,2}^{(2)}(\tau), \quad \chi_{2,2}^{Vir(3)}(\tau) = \eta(\tau)c_{1,1}^{(2)}(\tau), \tag{20}$$

and

$$\chi_{1,1}^{Vir(5)}(\tau) + \chi_{4,1}^{Vir(5)}(\tau) = \eta(\tau)c_{0,0}^{(3)}(\tau), \quad \chi_{4,3}^{Vir(5)}(\tau) = \eta(\tau)c_{3,1}^{(3)}(\tau) \tag{21}$$

$$\chi_{2,1}^{Vir(5)}(\tau) + \chi_{3,1}^{Vir(5)}(\tau) = \eta(\tau)c_{2,0}^{(3)}(\tau), \quad \chi_{3,3}^{Vir(5)}(\tau) = \eta(\tau)c_{1,1}^{(3)}(\tau). \tag{22}$$

The deep rôle of this representation in the description of noncritical $N = 2$ strings with minimal, unitary matter is yet to be uncovered. However, it is worth noticing that there is a direct relation between the parafermionic algebra involved in the representation of the $\hat{sl}(2|1; \mathbb{C})_k$ currents at level $k = \frac{1}{u} - 1$, and the parafermionic algebra involved in the realisation of the $N = 2$ superconformal algebra when the central charge is (2) with $p = 1$. The former is $\mathbb{Z}_{u-1}$ while the latter is $\mathbb{Z}_{u-2}$.

Acknowledgements

We would like to thank H. Kausch for useful comments. M. Hayes acknowledges the British EPSRC for a studentship. A. Taormina acknowledges the Leverhulme Trust for a fellowship and thanks the EC for support under a Training and Mobility of Researchers Grant No. FMRX-CT-96-0012.

References

- [1] Aharony, G., Ganor, O., Sonnenschein, J., Yankielowicz, S., Sochen, N., Nucl. Phys. B**399** (1993) 527; Phys. Lett. B**289** (1992) 309; Phys. Rev. Lett. **66** (1991) 2051.
- [2] Fan, J-B., Yu, M., ‘*Modules over affine Lie superalgebras*’, AS-ITP-93-14 ; ‘*G/G gauged supergroup valued WZWN field theory*’, hep-th 9304122.
- [3] Hu, H.L., Yu, M., Phys. Lett. B**289** (1992) 302; Nucl. Phys. B**391** (1993) 389.
- [4] Kac, V.G., Wakimoto M., Proc. Nat. Acad. Sci. **85** (1988) 4956.
- [5] Hayes M.R. and Taormina A., ‘*Admissible $SL(2/1)$ characters and parafermions*’, DTP 98/7, hep-th 9803022.
- [6] Bowcock P. and Taormina A. Commun. Math. Phys.**185** (1997) 467.
- [7] Bowcock P., Koktava R-L. K. and Taormina A., Phys. Lett. B**388** (1996) 303.
- [8] Bowcock P., Hayes M.R. and Taormina A., Nucl. Phys. B**510** [PM] (1998) 739.
- [9] Kac V.G. and Peterson D.H., Adv. in Math. **53** (1984) 125.
- [10] Fateev V.A. and Zamolodchikov A.B., Sov. Phys. JETP **62** (2) (1985) 215 ; Sov. Phys. JETP **63** (5) (1986) 913.
- [11] Qiu Z., Phys. Lett. B**198** (1987) 497.
- [12] Gepner D. and Qiu Z., Nucl. Phys. B**285** (1987) 423.
- [13] Dijkgraaf R., Vafa C., Verlinde E. and Verlinde H., Commun. Math. Phys.**123** (1989) 485.
- [14] Awata H. and Yamada Y., Mod. Phys. Lett A**7** (1992) 1185.
- [15] Huitu K., Nemeshansky D. and Yankielowicz S., Phys. Lett. B**246** (1990) 105.